

TIM-4 antibody [RMT4-54] (PE)

Cat. No. GTX01497-08

宿主	Rat
克隆	Monoclonal
同种型	IgG2a
实验应用	FCM
种属反应	Mouse

引用文献 (1)

产品说明

摘要

The RMT4-54 antibody reacts with mouse Tim-4, also known as T cell immunoglobulin and mucin domain containing protein-4. Tim-4 is expressed on macrophages and dendritic cells and can serve as a receptor for Tim-1. Tim-4 also functions as a receptor for phosphatidylserine and serves to enhance the phagocytosis of apoptotic cells.

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution

Recommended dilution

FCM

Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

属性


形式	Liquid
存储溶液	10mM NaH ₂ PO ₄ , 150mM NaCl, 0.1% Gelatin
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.
浓度	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Phycoerythrin (PE) Wavelength
注意事项	仅供实验室使用。不适用于人类或动物的任何临床，治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).

Date 2026 / 04 / 18 Page 1 of 2

產品圖片

**GTX01497-08 FCM Image**

FACS analysis of mouse C57Bl/6 resident peritoneal exudate cells using GTX01497-08 TIM-4 antibody [RMT4-54] (PE).

Right panel : co-stained with TIM-4 antibody [RMT4-54] (PE) and Mouse CD11b antibody (APC)

Left panel : co-stained with isotype control and Mouse CD11b antibody (APC)

antibody amount : 0.125 μ g (5 μ l)



For full product information, images and publications, please visit our [website](#).