

CSK antibody

Cat. No. GTX107916

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, ICC/IF, IHC-P
种属反应	Human

引用文献 (2)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片


IHC-Fr: 冰冻切片

属性


形式	Liquid
存储溶液	0.1M Tris, 0.1M Glycine, 10% Glycerol
保存剂	0.01% Thimerosal
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_2036680
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。

For full product information, images and publications, please visit our [website](#).


產品圖片

**GTX107916 WB Image**


Wild-type (WT) and CSK knockout (KO) 293T cell extracts (30 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with CSK antibody (GTX107916) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

**GTX107916 ICC/IF Image**

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using c-Src(GTX107916) antibody at 1:200 dilution.

**GTX107916 WB Image**

Sample (30 μ g whole cell lysate)
 A:293T
 B:H1299
 C:MOLT4 (GTX27912)
 7.5% SDS PAGE
 GTX107916 diluted at 1:1000

**GTX107916 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded HT29 xenograft , using c-Src(GTX107916) antibody at 1:100 dilution.
 Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](#).