

cIAP2 antibody [C1C3]

Cat. No. GTX113128

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应 用	WB, IHC-P
种属反应	Human, Rat

引用文献(2)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

以下为常规应用缩写的中文注解

WB: **免疫印迹** ICC/IF: 细**胞染色** IHC-P: **石蜡切片** IHC-Fr: 冰冻**切片**

形式 Liquid 存储溶液 PBS, 20% Glycerol 保存剂 0.025% ProClin 300 存放说明 Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
保存剂 0.025% ProClin 300 Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For
Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For
左放 说船
浓度 0.67 mg/ml (Please refer to the vial label for the specific concentration.)
偶 联 Unconjugated
RRID AB_2036572
注意事项




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 07 Page 1 of 2



產品图片



GTX113128 WB Image


cIAP2 antibody detects cIAP2 protein by western blot analysis.

A. 50 µg rat spleen lysate/extract

7.5% SDS-PAGE

cIAP2 antibody (GTX113128) dilution: 1:500


The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX113128 IHC-P Image


Immunohistochemical analysis of paraffin-embedded SW480 xenograft, using cIAP2(GTX113128) antibody at 1:100 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX113128 WB Image

cIAP2 antibody detects cIAP2 protein by western blot analysis. HeLa whole cell extract, HeLa cytosol and HeLa nuclear extract (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with cIAP2 antibody (GTX113128) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX113128 WB Image

Various whole cell extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with cIAP2 antibody [C1C3] (GTX113128) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 07 Page 2 of 2