

BRD3 antibody

Cat. No. GTX115058

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应 用	WB, ICC/IF, IHC-P
种属反应	Human, Mouse

引用文献(1)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent
以下为常规应用缩写的中文注解	

WB: 免疫印迹 ICC/IF: 细胞染色 IHC-P: 石蜡切片 IHC-Fr: 冰冻切片


属性	
形式	Liquid
存储溶液	PBS, 20% Glycerol
保存 剂	0.025% ProClin 300
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓 度	0.54 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_10724114
注意事项	仅 供 实验 室使用。不适用于人类或动物的任何临床,治疗或诊断用途。不适合 动 物或人 类 食用。



For full product information, images and publications, please visit our <u>website</u>.


Date 2025 / 06 / 04 Page 1 of 2

產品图片




GTX115058 WB Image

HeLa whole cell and nuclear extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with BRD3 antibody (GTX115058) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX115058 ICC/IF Image


BRD3 antibody detects BRD3 protein at nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: BRD3 stained by BRD3 antibody (GTX115058) diluted at 1:500. Red: phalloidin, a cytoskeleton marker, diluted at 1:200. Scale bar= $10 \ \mu m$.



GTX115058 WB Image

Sample (20 µg of whole cell lysate) A: mouse Liver 7.5% SDS PAGE GTX115058 diluted at 1:20000

 $\label{thm:conjugated} The \ HRP-conjugated \ anti-rabbit \ IgG \ antibody \ (GTX213110-01) \ was \ used \ to \ detect \ the \ primary \ antibody.$



GTX115058 IHC-P Image

BRD3 antibody detects BRD3 protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded mouse testis.
BRD3 stained by BRD3 antibody (GTX115058) diluted at 1:500.
Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 04 Page 2 of 2