

GITR antibody [DTA-1]

Cat. No. GTX14350

宿主	Rat
克隆	Monoclonal
同种型	IgG2b
实验应用	Activation
种属反应	Mouse

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
Activation	Assay dependent

Note : The DTA-1 antibody is an agonistic antibody that is commonly used to induce GITR signaling in vivo.

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

属性

形式	Liquid
存储溶液	Filter-sterilized PBS
保存剂	No preservative
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
浓度	Batch dependent (Please refer to the vial label for the specific concentration.)
纯度	>95% (Determined by SDS-PAGE)
内毒素	< 0.002 EU/μg (Determined by LAL assay)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).