

## alpha 1,2 Mannosidase IA antibody

**Cat. No. GTX40044**

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, ICC/IF, IHC-P
种属反应	Human, Mouse, Rat

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	0.5-1 µg/ml
ICC/IF	5-10 µg/ml
IHC-P	Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


## 属性

形式	Liquid
存储溶液	PBS, 1% BSA
保存剂	15mM Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	Batch dependent (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).


## 產品圖片

**GTX40044 ICC/IF Image**


ICC/IF analysis of NRK cells using GTX40044 alpha 1,2 Mannosidase IA antibody at 10 µg/mL. Cells were fixed and permeabilized with cold methanol followed by cold methanol:acetone (1:1).

**GTX40044 IHC-P Image**

IHC-P analysis of mouse testis tissue using GTX40044 alpha 1,2 Mannosidase IA antibody at 0.4 µg/mL.

**GTX40044 IHC-P Image**

IHC-P analysis of rat kidney tissue using GTX40044 alpha 1,2 Mannosidase IA antibody at 0.2 µg/mL.

**GTX40044 WB Image**

WB analysis of NIH3T3 cells using GTX40044 alpha 1,2 Mannosidase IA antibody.

Lane 1 : 1 µg/mL antibody

Lane 2 : 1 µg/mL antibody + immunizing peptide



For full product information, images and publications, please visit our [website](#).