

CPS1 antibody, Internal

Cat. No. GTX46535

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, IHC-P
种属反应	Human, Pig

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	Assay dependent
IHC-P	2-10 ug/ml

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	PBS, 2% Sucrose
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).

產品圖片




GTx46535 IHC-P Image

IHC-P analysis of Pig duodenum tissue using GTx46535 CPS1 antibody at 1:500.



GTx46535 WB Image

WB analysis of human fetal liver tissue using GTx46535 CPS1 antibody at 5.0µg/ml.



GTx46535 IHC-P Image

IHC-P analysis of human intestine tissue using GTx46535 CPS1 antibody at 4.0-8.0µg/ml.



GTx46535 IHC-P Image

IHC-P analysis of Pig ileum tissue using GTx46535 CPS1 antibody at 1:500.



GTx46535 IHC-P Image

IHC-P analysis of Pig kidney tissue using GTx46535 CPS1 antibody at 1:500.



For full product information, images and publications, please visit our [website](#).