

C1q antibody [JL-1] (FITC)

Cat. No. GTX54404-06

宿主	Mouse
克隆	Monoclonal
同种型	IgG2b
实验应用	WB, ICC/IF, IHC-Fr, ELISA, Depletion
种属反应	Human, Mouse, Rat

引用文献 (3)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	The typical starting working dilution is 1:50.
ICC/IF	The typical starting working dilution is 1:50.
IHC-Fr	The typical starting working dilution is 1:50.
ELISA	Assay dependent
Depletion	The typical starting working dilution is 1:50.

Note : Antibody JL-1 was used to stain tissue sections which were fixed in acetone.**Antibody JL-1 was administered to mice resulting in depletion of circulating C1q, glomerular deposition of C1q and induction of anti-C1q autoantibodies in susceptible mice.**

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

属性

形式	Liquid
存储溶液	Filter-sterilized PBS, 1% BSA
保存剂	0.02% sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light.
浓度	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Fluorescein isothiocyanate (FITC) Wavelength
注意事项	仅供实验室使用。不适用于人类或动物的任何临床，治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).