

E-Cadherin antibody [HL1229]

Cat. No. GTX636577

| 宿主 | Rabbit |
|--------------|-----------------------------|
| 克隆 | Monoclonal |
| 同种型 | IgG |
| 实验应 用 | WB, ICC/IF, IHC-P |
| 种属反应 | Human, Mouse, Rat, Cat, Dog |

引用文献(3)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:3000 |
| ICC/IF | Assay dependent |
| IHC-P | 1:100-1:1000 |

以下为常规应用缩写的中文注解

WB: 免疫印迹 ICC/IF: 细胞染色 IHC-P: 石蜡切片 IHC-Fr: 冰冻切片

| 属性 | |
|--------------|--|
| 形式 | Liquid |
| 存储溶液 | PBS |
| 保存 剂 | No preservative |
| 存放说明 | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| 浓度 | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| 偶联 | Unconjugated |
| RRID | AB_2910011 |
| 注意事 项 | 仅 供 实验 室使用。不适用于人类或动物的任何临床,治疗或诊断用途。不适合动物或人类食用。 |
| | |




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 14 Page 1 of 2




產品图片




GTX636577 WB Image

Non-transfected (–) and transfected (+) MCF-7 whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with E-Cadherin antibody [HL1229] (GTX636577) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX636577 WB Image

Whole cell extract (30 μ g) was separated by 5% SDS-PAGE, and the membrane was blotted with E-Cadherin antibody [HL1229] (GTX636577) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX636577 WB Image

Mouse tissue extract (50 μ g) was separated by 5% SDS-PAGE, and the membrane was blotted with E-Cadherin antibody [HL1229] (GTX636577) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX636577 IHC-P Image

E-Cadherin antibody [HL1229] detects E-Cadherin protein at cell membrane by immunohistochemical analysis.

Sample: Paraffin-embedded mouse intestine.

E-Cadherin stained by E-Cadherin antibody [HL1229] (GTX636577) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX636577 IHC-P Image

E-Cadherin antibody [HL1229] detects E-Cadherin protein at cell membrane by immunohistochemical analysis.

Sample: Paraffin-embedded mouse colon.

E-Cadherin stained by E-Cadherin antibody [HL1229] (GTX636577) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 14 Page 2 of 2