

MC1 Receptor antibody [HL1471]

Cat. No. GTX636945

宿主	Rabbit
克隆	Monoclonal
同种型	IgG
实验应用	WB, ICC/IF, IHC-P, IHC-Fr
种属反应	Human, Mouse, Rat

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	1:100-1:1000
IHC-Fr	Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	PBS
保存剂	No preservatives
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).


產品圖片



GTX636945 WB Image

Non-transfected (-) and transfected (+) unboiled A375 whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membranes were blotted with MC1 Receptor antibody [HL1471] (GTX636945) diluted at 1:5000 and competitor's antibody (Highly Cited Competitor) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


*The competitor is not affiliated with GeneTex and does not endorse this product.



GTX636945 WB Image

Various tissue extracts (50 µg) was separated by 12% SDS-PAGE, and the membranes were blotted with MC1 Receptor antibody [HL1471] (GTX636945) diluted at 1:1000 and competitor's antibody (Competitor) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.


*The competitor is not affiliated with GeneTex and does not endorse this product.



GTX636945 WB Image


Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membranes were blotted with MC1 Receptor antibody [HL1471] (GTX636945) diluted at 1:50000 and competitor's antibody diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

*The competitor is not affiliated with GeneTex and does not endorse this product.



GTX636945 WB Image

Various tissue extracts (50 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with MC1 Receptor antibody [HL1471] (GTX636945) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX636945 IHC-P Image

MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane by immunohistochemical analysis.

Sample: Paraffin-embedded A375 xenograft .

MC1 Receptor stained by MC1 Receptor antibody [HL1471] (GTX636945) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).