

U2AF65 antibody [HL1583]

Cat. No. GTX637062

宿主	Rabbit
克隆	Monoclonal
同种型	IgG
实验应用	WB, IHC-P
种属反应	Human, Mouse, Rat

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片


IHC-Fr: 冰冻切片

属性

形式	Liquid
存储溶液	PBS
保存剂	No preservative
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).


產品圖片

GTx637062 IHC-P Image

U2AF65 antibody [HL1583] detects U2AF65 protein by immunohistochemical analysis.

Sample: Paraffin-embedded rat tissue.

U2AF65 stained by U2AF65 antibody [HL1583] (GTx637062) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTx637062 IHC-P Image

U2AF65 antibody [HL1583] detects U2AF65 protein by immunohistochemical analysis.


Sample: Paraffin-embedded mouse tissue.

U2AF65 stained by U2AF65 antibody [HL1583] (GTx637062) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTx637062 WB Image

Various whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with U2AF65 antibody [HL1583] (GTx637062) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



GTx637062 IHC-P Image

U2AF65 antibody [HL1583] detects U2AF65 protein at nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded rat testis.

U2AF65 stained by U2AF65 antibody [HL1583] (GTx637062) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTx637062 IHC-P Image

U2AF65 antibody [HL1583] detects U2AF65 protein at nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded mouse liver.

U2AF65 stained by U2AF65 antibody [HL1583] (GTx637062) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](http://www.genetex.com).