

## Macrophage + Monocyte antibody [MAC387]

Cat. No. GTX76577

宿主	Mouse
克隆	Monoclonal
同种型	IgG1
实验应用	IHC-P, IHC-Fr, FCM
种属反应	Human, Mouse, Rat, Rabbit, Goat, Bovine, Cat, Dog, Guinea pig, Baboon, Cynomolgus monkey, Fallow deer, Horse, MarmosetPig, Mink, Pygmy hippopotamus, Rhesus Monkey, Squirrel monkey

引用文献 (6)

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
IHC-P	1/100-1/200
IHC-Fr	1/100-1/200
FCM	1/50-1/100

**Note : This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase. Permeabilization step is needed.**

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

## 属性

形式	Liquid
存储溶液	PBS
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_377426
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).