

MDM2 antibody, Internal

Cat. No. GTX89355

宿主	Goat
克隆	Polyclonal
同种型	IgG
实验应用	WB
种属反应	Human

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	0.3-1µg/ml

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片


IHC-Fr: 冰冻切片

属性


形式	Liquid
存储溶液	TBS, 0.5% BSA
保存剂	0.02% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).

產品圖片

GTX89355 WB Image

WB analysis of HEK293 overexpressing MDM2 (mock transfection in first lane) using GTX89355 MDM2 antibody, Internal.


GTX89355 WB Image

WB analysis of human liver lysate using GTX89355 MDM2 antibody, Internal.

Dilution : 0.3μg/ml

Loading : 35μg protein in RIPA buffer



For full product information, images and publications, please visit our [website](https://www.genetex.com).